arXiv:1301.1491v3 [math.KT] 6 May 2013

EQUIVARIANT ALGEBRAIC KK-THEORY AND ADJOINTNESS THEOREMS

EUGENIA ELLIS

Abstract. We introduce an equivariant algebraic kk-theory for G-algebras and G-graded algebras. We study some adjointness theorems related with crossed product, trivial action, induction and restriction. In particular we obtain an algebraic version of the Green-Julg Theorem which gives us a computational tool.

1. Introduction

Algebraic kk-theory has been introduced by G. Cortiñas and A. Thom in [2]. This is a bivariant K-theory on the category of ℓ-algebras where ℓ is a commutative ring with unit. For each pair (A, B) of ℓ-algebras a group $\mathrm{kk}(A, B)$ is defined. A category $\mathfrak{KK}$ is obtained whose objects are ℓ-algebras and where the morphisms from A to B are the elements of the group $\mathrm{kk}(A, B)$. The category $\mathfrak{KK}$ is triangulated and there is a canonical functor $j : \mathrm{Alg}_\ell \to \mathfrak{KK}$ with universal properties. These properties are algebraic homotopy invariance, matrix invariance and excision.

The definition of algebraic kk-theory was inspired by the work of J. Cuntz [4] and N. Higson [7] on the universal properties of Kasparov KK-theory [8]. The KK-theory of separable C^*-algebras is a common generalization both of topological K-homology and topological K-theory as an additive bivariant functor. Let A, B be separable C^*-algebras. Then

$$KK_*(\mathbb{C}, B) \simeq K_*^{top}(B) \qquad KK^*(A, \mathbb{C}) = K^*_{hom}(A) \tag{1.1}$$

here $K_*^{top}(B)$ denotes the K-theory of B and $K^*_{hom}(A)$ the topological K-homology of A. J. Cuntz in [3] gave another equivalent definition of the original one given in [8]. This new approach allowed to put bivariant K-theory in algebraic context. Higson in [7] stated the universal property of KK whose algebraic analogue is studied in [2], where also an analogue of (1.1) is proved. On the algebraic side, if A is an ℓ-algebra then

$$\mathrm{kk}(\ell, A) \simeq \mathrm{KH}(A)$$

here KH is Weibel's homotopy K-theory defined in [12]. We can start to build a dictionary between Kasparov's KK-theory and algebraic kk-theory in the following way

The author was partially supported by CSIC-Uruguay; by the MathAmSud network U11MATH-05 partially financed by ANII (Uruguay) and MINCyT (Argentina) and by grants UBACyT 20020100100386, and MTM2007-64704 (FEDER funds).

Kasparov's KK-theory	$\leftrightarrow$	algebraic kk-theory
bivariant K-theory on separable C^*-algebras C^*-Alg	$\leftrightarrow$	bivariant K-theory on ℓ-algebras Alg_ℓ
$k : C^*\text{-Alg} \to KK$	$\leftrightarrow$	$j : \mathrm{Alg}_\ell \to \mathfrak{RR}$
k is stable with respect to compact operators $\mathcal{K}$	$\leftrightarrow$	j is stable with respect to $M_\infty = \bigcup_{n\in\mathbb{N}} M_n$
k is continous homotopy invariant	$\leftrightarrow$	j is polynomial homotopy invariant
k is split exact	$\leftrightarrow$	j is excisive
k is universal for the properties described above	$\leftrightarrow$	j is universal for the properties described above
$\mathrm{KK}_*(\mathbb{C}, A) \simeq K_*^{top}(A)$	$\leftrightarrow$	$\mathrm{kk}_*(\ell, A) \simeq \mathrm{KH}_*(A)$

In this paper we obtain an equivariant version of the dictionary stated above in the following sense

Equivariant Kasparov's KK-theory	$\leftrightarrow$	Equivariant algebraic kk-theory
bivariant K-theory on separable G-C^*-algebras	$\leftrightarrow$	bivariant K-theory on G-algebras
$k : G\text{-}C^*\text{-Alg} \to KK^G$	$\leftrightarrow$	$j^G : G\text{-Alg} \to \mathfrak{RR}^G$
k is stable with respect to $\mathcal{K}(\ell^2(G \times \mathbb{N}))$	$\leftrightarrow$	j^G is G-stable
k is continous homotopy invariant	$\leftrightarrow$	j^G is polynomial homotopy invariant
k is split exact	$\leftrightarrow$	j^G is excisive
k is universal for the properties described above	$\leftrightarrow$	j^G is universal for the properties described above
$\mathrm{KK^G}_*(\mathbb{C}, A) \simeq K_*^{top}(A \rtimes G)$ with G compact	$\leftrightarrow$	$\mathrm{kk}^G(\ell, A) \simeq \mathrm{KH}(A \rtimes G)$ with G finite and $1/\lvert G\rvert \in \ell$

We also introduce a dual theory $\hat{\mathfrak{RR}}^G$ for G-graded algebras and establish a duality result similar to that proved by Baaj and Skandalis in [1].

In Section 2 we recall some results from [2]. We take special care in the definition of equivariant matrix invariance. We introduce the concept of G-stable functor in Section 3, the rest of the concepts which appears in the equivariant dictionary are straightforward. The definition of G-stability was inspired by the definition of equivariant stability for G-C^*-algebras (see [11]). In sections 2 and 3 G is a group without any other assumption but from Section 4 and for the rest of the paper G is a countable group.

In Section 4 we introduce the appropiate brand of equivariant algebraic kk-theory in each case and establish its universal properties. For a countable group G, we

define an equivariant algebraic kk-theory $\mathfrak{KK}^G$ for the category of G-algebras and $\hat{\mathfrak{KK}}^G$ for the category of G-graded algebras.

We study adjointness theorems in equivariant kk-theory. We put in an algebraic context some of the adjointness theorems which appear in Kasparov KK-theory. In Section 5, we define the functors of trivial action and crossed product between $\mathfrak{KK}$ and $\mathfrak{KK}^G$. The first adjointness theorem is Theorem 5.2.1 which is an algebraic version of the Green-Julg Theorem. This result gives us the first computation related with homotopy K-theory. If G is a finite group, A is a G-algebra, B is an algebra and $\frac{1}{|G|} \in \ell$ then there is an isomorphism

$$\psi_{GJ} : \mathrm{kk}^G(B^\tau, A) \to \mathrm{kk}(B, A \rtimes G).$$

In particular, if $B = \ell$ then

$$\mathrm{kk}^G(\ell, A) \simeq \mathrm{KH}(A \rtimes G).$$

In Section 6 we consider H a subgroup of G. We define induction and restriction funtors between $\mathfrak{KK}^G$ and $\mathfrak{KK}^H$ and study the adjointness between them. If B is an H-algebra and A is a G-algebra then there is an isomorphism

$$\psi_{IR} : \mathrm{kk}^G(\mathrm{Ind}_{\mathrm{H}}^{\mathrm{G}} B, A) \to \mathrm{kk}^H(B, \mathrm{Res}_{\mathrm{G}}^{\mathrm{H}} A).$$

This result gives us another computation. Taking H the trivial group and $B = \ell$ we obtain that

$$\mathrm{kk}^G(\ell^{(G)}, A) \simeq \mathrm{KH}(A) \qquad \forall A \in G - \mathrm{Alg}\,.$$

Here $\ell^{(G)} = \bigoplus_{g \in G} \ell$ with the regular action of G. More general, if H is a finite subgroup of G and $1/|H| \in \ell$ we combine ψ_{GJ} and ψ_{IR} and obtain

$$\mathrm{kk}^G(\ell^{(G/H)}, A) \simeq \mathrm{KH}(A \rtimes H) \qquad \forall A \in G - \mathrm{Alg}\,.$$

In Section 7 we obtain an algebraic version of the Baaj-Skandalis duality theorem. We show that the functors

$$\rtimes G : \mathfrak{KK}^G \to \hat{\mathfrak{KK}}^G \qquad G \hat{\rtimes} : \hat{\mathfrak{KK}}^G \to \mathfrak{KK}^G$$

are inverse category equivalences.

Acknowledgements. The results here are part of my PhD Thesis. I am thankful to Willie Cortiñas for his orientation and comments about this paper. I would like also to thank the referee for useful comments. This paper was finished during a visit to the Freie Universität Berlin (FU), financed by Berlin Mathematical School (BMS) and International Mathematical Union (IMU). I am also thankful to Freie Universität for its hospitality and to BMS-IMU for its support.

2. Algebraic kk-theory

In this section we recall some results from [2] and we adapt them to our setting. Let ℓ be a commutative ring with unit. We consider an ℓ-bimodule A such that $x \cdot a = a \cdot x$, with $x \in \ell$ and $a \in A$. An ℓ-bimodule A is an ℓ-algebra if it is an associative and not necessarily unital algebra. Let G be a group. A G-algebra A is an ℓ-algebra with an action of G, i.e. with a group homomorphism $\alpha : G \to \mathrm{End}_\ell(A)$, where $\mathrm{End}_\ell(A)$ denote the group of ℓ-linear endomorphisms of A. We shall denote by $g(a)$ or $g \cdot a$ the element $\alpha(g)(a)$. An equivariant morphism $f : A \to B$ between

G-algebras is a G-equivariant ℓ-linear map. A G-graded algebra A is an ℓ-algebra with a familiy of ℓ-submodules $\{A_s\}_{s\in G}$ such that

$$A = \bigoplus_{s\in G} A_s \qquad\qquad A_sA_t \subseteq A_{st} \qquad\qquad s,t \in G.$$

We write $|a| = s$ if $a \in A_s$. An homogeneous morphism $f : A \to B$ of G-graded algebras is an algebra morphism such that $f(A_s) \subset B_s$, for all $s \in G$. We consider the category of G-algebras with equivariant morphisms and the category of G-graded algebras with homogeneous morphisms.

In this section we define an algebraic kk-theory for the categories of G-algebras and G-graded algebras. We write $\mathcal{C}$ to refer to either of these categories.

2.1. **Homotopy invariance.** Let A be an object of $\mathcal{C}$. Put $A^{\Delta^1} := A[t] = A \otimes_{\mathbb{Z}} \mathbb{Z}[t]$. Consider the trivial action of G (trivial G-grading) on $\mathbb{Z}[t]$ and the diagonal structure on $A[t]$. Then $A[t]$ is an object of $\mathcal{C}$. Let us write $c_A : A \to A[t]$ for the inclusion of A as constant polynomials in $A[t]$ and $\mathrm{ev}_i : A[t] \to A$ for the evaluation of t at i ($i = 0, 1$). Note that these maps are morphisms in $\mathcal{C}$ and that c_A is a section of ev_i.

Let $f_0, f_1 : A \to B$ be morphisms in $\mathcal{C}$. We call f_0 and f_1 elementarily homotopic, and write $f_0 \sim_e f_1$, if there exists a morphism $H : A \to B[t]$ such that $\mathrm{ev}_i H = f_i$, $i = 0, 1$. It is easy to check that elementary homotopy is a reflexive and symmetric relation but in general it is not transitive. Let $f, g : A \to B$ be morphisms in $\mathcal{C}$. We call f and g homotopic, and write $f \sim g$, if they can be connected by a finite chain of elementary homotopies,

$$f \sim_e h_0 \sim_e \ldots \sim_e h_n \sim_e g.$$

We denote the set of homotopy clases by $[A, B]_{\mathcal{C}}$. A morphism $f : A \to B$ is an elementary homotopy equivalence if there exists a morphism $g : B \to A$ such that $f \circ g \sim_e \mathrm{id}_B$ and $g \circ f \sim_e \mathrm{id}_A$. We say A is elementarily contractible if the null morphism and the identity morphism are elementarily homotopic.

The category of ind-objects of $\mathcal{C}$ is the category ind-$\mathcal{C}$ of directed diagrams in $\mathcal{C}$. An object in ind-$\mathcal{C}$ is described by a filtering partially ordered set $(I, \leq)$ and a functor $A : I \to \mathcal{C}$. The set of homomorphisms in ind-$\mathcal{C}$ is defined by

$$\hom_{\text{ind-}\mathcal{C}}((A, I), (B, J)) := \lim_{i\in I} \operatorname*{colim}_{j\in J} \hom_{\mathcal{C}}(A_i, B_j).$$

Let $A = (A, I)$ and $B = (B, J)$ be objects of ind-$\mathcal{C}$, we have a map

$$\hom_{\text{ind-}\mathcal{C}}(A, B) \mapsto [A, B]_{\mathcal{C}} = \lim_{i\in I} \operatorname*{colim}_{j\in J} [A_i, B_j]_{\mathcal{C}}. \tag{2.1.1}$$

We say two morphisms in ind-$\mathcal{C}$ are homotopic if their images by (2.1.1) are equal. An object of ind-$\mathcal{C}$ is contractible if the null morphism and the identity morphism are homotopic. Sometimes we shall omit the poset in the notation.

Let $\mathcal{D}$ be an arbitrary category. A functor $F : \mathcal{C} \to \mathcal{D}$ is homotopy invariant if it maps the inclusion $c_A : A \to A[t]$ to an isomorphism. It is easy to check that $F : \mathcal{C} \to \mathcal{D}$ is an homotopy invariant functor if and only if $F(f) = F(g)$ when $f \sim g$ (or equivalenty when $f \sim_e g$). If $\mathcal{C}$ is the category of G-algebras we say that an homotopy invariant functor is equivariantly homotopy invariant and if $\mathcal{C}$ is the category of G-graded algebras we say that a homotopy invariant functor is graded homotopy invariant.

2.2. **Matrix stability.** Consider M_n the algebra of $n\times n$-matrices with coefficients in $\mathbb{Z}$ with the trivial action (grading) of G and $M_\infty = \cup_{n\geq 0} M_n$. Let A be an object in $\mathcal{C}$. We define

$$M_n A = M_n \otimes_{\mathbb{Z}} A \qquad M_\infty A = M_\infty \otimes_{\mathbb{Z}} A$$

which are objects in $\mathcal{C}$ with the diagonal action (grading). Denote by $\iota_n : \mathbb{Z} \to M_n$ and $\iota_\infty : \mathbb{Z} \to M_\infty$ the inclusions at the upper left corner. A functor $F : \mathcal{C} \to \mathcal{D}$ is M_n-stable (M_∞-stable) if $F(\iota_n \otimes \mathrm{id}_A)$ ($F(\iota_\infty \otimes \mathrm{id}_A)$) is an isomorphism for all $A \in \mathcal{C}$.

2.3. **Algebra of polynomial functions.** Consider the following simplicial ring

$$\mathbb{Z}^\Delta : [n] \mapsto \mathbb{Z}^{\Delta^n} \qquad \mathbb{Z}^{\Delta^n} := \mathbb{Z}[t_0, \ldots, t_n]/ < 1 - \sum_i t_i > \tag{2.3.1}$$

$$\begin{array}{rcll} \Theta : [n] \to [m] & \mapsto & \Theta^* : \mathbb{Z}^{\Delta^m} \to \mathbb{Z}^{\Delta^n} & \\ & & \Theta^*(t_i) = \left\{ \begin{array}{ll} 0 & \text{si } \Theta^{-1}(i) = \emptyset \\ \sum_{j\in\Theta^{-1}(i)} t_j & \text{si } \Theta^{-1}(i) \neq \emptyset \end{array} \right. & \end{array}$$

Let A be an object in $\mathcal{C}$. Define

$$A^\Delta : [n] \mapsto A^{\Delta^n} \qquad A^{\Delta^n} := A \otimes_{\mathbb{Z}} \mathbb{Z}^{\Delta^n}$$

Note A^Δ is a simplicial $\mathcal{C}$-object. Let X be a simplicial set. Define

$$A^X := \mathrm{map}_{\mathbb{S}}(X, A^\Delta)$$

If $(K, \star)$ is a pointed simplicial set, put

$$A^{(K,\star)} := \mathrm{map}_{\mathbb{S}_*}((K,\star), A^\Delta) = \ker(\mathrm{map}_{\mathbb{S}}(K, A^\Delta) \to \mathrm{map}_{\mathbb{S}}(\star, A^\Delta)) = \ker(A^K \to A)$$

If A is a G-algebra (G-graded algebra) and K a finite simplicial set, by [2, Lemma 3.1.3] we can consider A^K and $A^{(K,\star)}$ as a G-algebras (G-graded algebras) taking the diagonal structure with the trivial structure in $\mathbb{Z}^K$ and $\mathbb{Z}^{(K,\star)}$.

We will denote by $\mathrm{sd}^\bullet X$ the following pro-simplicial set

$$\mathrm{sd}^\bullet X : \ldots \to \mathrm{sd}^n X \xrightarrow{h} \mathrm{sd}^{n-1} X \to \ldots \to \mathrm{sd} X \xrightarrow{h} X$$

where $\mathrm{sd}X$ is the subdivision of X and h is the last vertex map, see [5, III.4]. If A is an object in $\mathcal{C}$ we consider the following object in ind-$\mathcal{C}$

$$A^{\mathrm{sd}^\bullet X} : A^X \to A^{\mathrm{sd}X} \to \ldots \to A^{\mathrm{sd}^{n-1}X} \to A^{\mathrm{sd}^n X} \to \ldots$$

2.4. **Extensions and classifying map.** A sequence of morphisms in ind-$\mathcal{C}$

$$A \xrightarrow{f} B \xrightarrow{g} C \tag{2.4.1}$$

is called an extension if f is a kernel of g and g is a cokernel of f. Let $\mathcal{U}(\mathcal{C})$ be the category of modules with linear and equivariant maps (graded maps), i.e. we forget the multiplication in $\mathcal{C}$ keeping the module structure. Let $F : \mathcal{C} \to \mathcal{U}(\mathcal{C})$ be the forgetful functor. This functor can be extended to $F : \text{ind-}\mathcal{C} \to \text{ind-}\mathcal{U}(\mathcal{C})$. We will call an extension (2.4.1) weakly split if $F(g)$ has a section in ind-$\mathcal{U}(\mathcal{C})$.

Let M be an object in $\mathcal{U}(\mathcal{C})$. Consider in

$$\tilde{T}(M) = \bigoplus_{n\geq 1} M^{\otimes^n} \quad M^{\otimes^n} = \underbrace{M \otimes \ldots \otimes M}_{n\text{-times}}$$

the usual structure of ℓ-algebra. If $\mathcal{C}$ is the category of G-algebras, we consider in $M^{\otimes^n}$ the following action,

$$g \cdot (m_1 \otimes m_2 \otimes \ldots \otimes m_n) = g \cdot m_1 \otimes g \cdot m_2 \otimes \ldots \otimes g \cdot m_n$$

which gives to $\tilde{T}(M)$ a G-algebra structure. If $\mathcal{C}$ is the category of G-graded algebras, take in $M^{\otimes^n}$ the following G-graduation

$$|m_1 \otimes m_2 \otimes \ldots \otimes m_n| = |m_1||m_2| \ldots |m_n|.$$

By this way $\tilde{T}(M)$ is a G-graded algebra. Both constructions are functorial hence we consider the functor $\tilde{T} : \mathcal{U}(\mathcal{C}) \to \mathcal{C}$. Put

$$T := \tilde{T} \circ F : \mathcal{C} \to \mathcal{C}$$

If A is an object in $\mathcal{C}$ there exists an morphism in $\mathcal{C}$

$$\eta_A : T(A) \to A \qquad \eta_A(a_1 \otimes \ldots \otimes a_n) = a_1 \ldots a_n$$

and a morphism in $\mathcal{U}(\mathcal{C})$ $\mu_A : A \to T(A)$ which is the inclusion at the first summand of $T(A)$. Let A, B be objects in $\mathcal{C}$, it is easy to check that

$$\hom_{\mathcal{C}}(T(A), B) \simeq \hom_{\mathcal{U}(\mathcal{C})}(F(A), F(B)). \tag{2.4.2}$$

Hence if we have a morphism $A \to B$ in $\mathcal{U}(\mathcal{C})$, we can extend it to a morphism $T(A) \to B$ in $\mathcal{C}$. It shows that T is the left adjoint of F. The counit of the adjuntion is $\eta_A : T(A) \to A$ and it is surjective (see [10, IV.3 Thm 1]). We define

$$J(A) := \ker \eta_A.$$

The universal extension of A is

$$J(A) \xrightarrow{\imath_A} T(A) \xrightarrow{\eta_A} A.$$

Let $A \xrightarrow{f} B \xrightarrow{g} C$ be a weakly split extension. Let s be a section of $F(g)$ and define $\hat{\xi} = \eta_B \circ \tilde{T}(s)$, then

$$\eta_C = \eta_C \circ T(g) \circ \tilde{T}(s) = g \circ \eta_B \circ \tilde{T}(s) = g \circ \hat{\xi}.$$

Define $\xi : J(C) \to A$ as the restriction of $\hat{\xi}$ to $J(C)$ We obtain a commutative diagram of extensions

$$\begin{array}{ccccc} A & \xrightarrow{f} & B & \xrightarrow{g} & C \\ \uparrow{\scriptstyle \xi} & & \uparrow{\scriptstyle \hat{\xi}} & & \uparrow{\scriptstyle \mathrm{id}_C} \\ J(C) & \xrightarrow[\imath_C]{} & T(C) & \xrightarrow[\eta_C]{} & C \end{array}$$

This construction of ξ depends on which s is chosen. If $\xi_1, \xi_2 : J(C) \to A$ are morphisms constructed taking different sections of $F(g)$ we will show ξ_1, ξ_2 are homotopic. Define a linear equivariant (graded) map

$$H : C \to A[t] \quad H(c) = (1-t)\xi_1(c) + t\xi_2(c).$$

By (2.4.2) it extends to a homomorphism and there exists a morphism $H : T(C) \to A[t]$ in $\mathcal{C}$ such that

$$\mathrm{ev}_0 \circ H_{|J(C)} = \xi_1 \qquad \mathrm{ev}_1 \circ H_{|J(C)} = \xi_2.$$

Then the map ξ is unique up to elementary homotopy. We call ξ the classifying map of the extension $A \xrightarrow{f} B \xrightarrow{g} C$. This construction is functorial, that means if we have a diagram of weakly split extensions,

$$\begin{array}{ccccc} A & \longrightarrow & B & \longrightarrow & C \\ {\scriptstyle f}\downarrow & & \downarrow{\scriptstyle h} & & \downarrow{\scriptstyle g} \\ A' & \longrightarrow & B' & \longrightarrow & C' \end{array}$$

then there is a diagram

$$\begin{array}{ccc} J(C) & \longrightarrow & A \\ {\scriptstyle J(g)}\downarrow & & \downarrow{\scriptstyle f} \\ J(C') & \longrightarrow & A' \end{array}$$

of classifying maps, which is commutative up to elementary homotopy.

Let L be a ring and A an object in $\mathcal{C}$. Then the extension

$$J(A) \otimes_{\mathbb{Z}} L \to T(A) \otimes_{\mathbb{Z}} L \to A \otimes_{\mathbb{Z}} L \tag{2.4.3}$$

is weakly split, and there is a choice for classifying map

$$\phi_{A,L} : J(A \otimes_{\mathbb{Z}} L) \to J(A) \otimes_{\mathbb{Z}} L$$

of (2.4.3), which is natural in both variables. In particular, if K is a finite pointed simplicial set. There is a homotopy class of maps

$$\phi_{A,K} : J(A^K) \to J(A)^K \tag{2.4.4}$$

natural with respect to K, which is represented by a classifying map of the following extension

$$J(A)^K \xrightarrow{\iota_A^*} T(A)^K \xrightarrow{\eta_A^*} A^K$$

Let S^1 be the simplicial circle $\Delta^1/\partial\Delta^1$, we define

$$\Omega := \mathbb{Z}^{(S^1,\star)} \qquad P := \mathbb{Z}^{(\Delta^1,\star)}$$

The path extension of A is the extension induced by the cofibration $\partial\Delta^1 \subset \Delta^1$ (see [2, Lemma 3.1.2]),

$$\Omega A \longrightarrow PA \xrightarrow{(\mathrm{ev}_0,\mathrm{ev}_1)} A \oplus A \tag{2.4.5}$$

The extension (2.4.5) is weakly split because we have a linear and equivariant (graded) section of $(\mathrm{ev}_0, \mathrm{ev}_1)$

$$(a, b) \mapsto (1 - t)a + tb. \tag{2.4.6}$$

The loop extension of A is

$$\Omega A \to PA \xrightarrow{\mathrm{ev}_1} A \tag{2.4.7}$$

Note $a \mapsto at$ is a natural section of $F(\mathrm{ev}_1)$. Thus we can pick a natural choice for the classifying map of (2.4.7). We call it

$$\rho_A : J(A) \to \Omega A$$

Let $\mathcal{P} := \mathbb{Z}^{(\mathrm{sd}^\bullet\Delta^1,\star)}$ we have an extension in ind-$\mathcal{C}$

$$A^{\mathcal{S}^1} := A^{(\mathrm{sd}^\bullet S^1,\star)} \qquad A^{\mathcal{S}^1} \to \mathcal{P}A \xrightarrow{\mathrm{ev}_1} A$$

which is naturally weakly split. The classifying map $J(A) \to A^{\mathcal{S}^1}$ is the following composition

$$J(A) \xrightarrow{\rho_A} \Omega A \xrightarrow{h} A^{\mathcal{S}^1} \tag{2.4.8}$$

where h is induced by the last vertex map. We will sometimes abuse notation and write ρ_A for the map (2.4.8).

Let $f : A \to B$ be a morphism in $\mathcal{C}$. The mapping path extension of f is the extension obtained from the loop extension of B by pulling it back to A

$$P_f := PB \times_B A \qquad \begin{array}{ccccc} \Omega B & \xrightarrow{\iota} & PB \times_B A & \xrightarrow{\pi_f} & A \\ \downarrow & & \downarrow & & \downarrow f \\ \Omega B & \xrightarrow[\iota]{} & PB & \xrightarrow[\mathrm{ev}_1]{} & B \end{array} \qquad \begin{array}{c} \mathrm{E}' \\ \\ \mathrm{E} \end{array}$$

We call P_f the path algebra of f. Note the extension E' is naturally weakly split because $\tilde{s}(a) = (s \circ f(a), a)$ is a natural section of π_f where s is the natural section of E. We define

$$\mathcal{P}_f := \mathcal{P}B \times_B A \qquad B^{\mathcal{S}^1} \to \mathcal{P}B \times_B A \xrightarrow{\pi_f} A. \tag{2.4.9}$$

Note $\rho_f := \rho_B J(f)$ is the classifying map of the extension (2.4.9).

2.5. **Excisive homology theories.** We consider triangulated categories in terms of a loop functor Ω. A triangulated category $(\mathcal{T}, \Omega, \mathcal{Q})$ is an additive category $\mathcal{T}$ with an equivalence $\Omega : \mathcal{T} \to \mathcal{T}$ and a class $\mathcal{Q}$ of sequences in $\mathcal{T}$ called distinguished triangles

$$(T) \qquad \Omega C \to A \to B \to C$$

satisfying some axioms, see [9], [2]. A triangle functor from $(\mathcal{T}_1, \Omega_1, \mathcal{Q}_1)$ to $(\mathcal{T}_2, \Omega_2, \mathcal{Q}_2)$ is a pair consisting of an additive functor $R : \mathcal{T}_1 \to \mathcal{T}_2$ and a natural transformation $\alpha : \Omega_2 R \to R\Omega_1$ such that

$$\Omega_2 R(C) \xrightarrow{R(f)\circ\alpha_C} R(A) \xrightarrow{R(g)} R(B) \xrightarrow{R(h)} R(C)$$

is a distinguished triangle in $\mathcal{T}_2$ for each triangle $\Omega_1 C \xrightarrow{f} A \xrightarrow{g} B \xrightarrow{h} C$.

Let $(\mathcal{T}, \Omega, \mathcal{Q})$ be a triangulated category. An excisive homology theory for $\mathcal{C}$ with values in $\mathcal{T}$ is an $\mathcal{E}$-excisive homology, as defined in [2, Sec 6.6], when $\mathcal{E}$ is the class of weakly split extensions.

Let P be a property for functors defined on $\mathcal{C}$ to some triangulated category. A functor $u : \mathcal{C} \to \mathcal{V}$ is a universal functor with P if it has the property P and if $F : \mathcal{C} \to \mathcal{T}$ is another functor with P there exists a unique triangle functor $G : \mathcal{V} \to \mathcal{T}$ such that the following diagram is commutative

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{u} & \mathcal{V} \\ & \searrow^{F} & \downarrow G \\ & & \mathcal{T} \end{array}$$

2.6. **The category $\mathfrak{KK}_{\mathcal{C}}$.** Let A and B be objects in $\mathcal{C}$. Consider $\mathcal{M}_\infty$ the ind-ring defined in [2, Sec 4.1] and define inductively $B^{\mathcal{S}^{n+1}} := (B^{\mathcal{S}^n})^{\mathcal{S}^1}$. Define

$$E_n(A,B)_{\mathcal{C}} := [J^n(A), \mathcal{M}_\infty B^{\mathcal{S}^n}]_{\mathcal{C}}.$$

Consider the following morphism $\imath_n : E_n(A,B)_{\mathcal{C}} \to E_{n+1}(A,B)_{\mathcal{C}}$ such that

$$\begin{array}{ccc} J^n(A) & \xrightarrow{f} & \mathcal{M}_\infty B^{\mathcal{S}^n} \\ & & \downarrow \\ J^{n+1}(A) & \xrightarrow{J(f)} & J(\mathcal{M}_\infty B^{\mathcal{S}^n}) \xrightarrow{\phi_{\mathcal{M}_\infty, B^{\mathcal{S}^n}}} \mathcal{M}_\infty J(B^{\mathcal{S}^n}) \xrightarrow{\rho_{B^{\mathcal{S}^n}}} \mathcal{M}_\infty B^{\mathcal{S}^{n+1}} \end{array}$$

Define

$$\mathrm{kk}_{\mathcal{C}}(A,B) = \mathrm{colim}_{n\in\mathbb{N}} E_n(A,B)_{\mathcal{C}}.$$

Let A, B and C be objects in $\mathcal{C}$. There exists an associative product

$$\circ : \mathrm{kk}_{\mathcal{C}}(B,C) \times \mathrm{kk}_{\mathcal{C}}(A,B) \to \mathrm{kk}_{\mathcal{C}}(A,C)$$

which extends the composition of algebra homomorphisms. If $[\alpha] \in \mathrm{kk}_{\mathcal{C}}(B,C)$ is an element represented by $\alpha : J^n(B) \to C^{\mathcal{S}^n}$ and $[\beta] \in \mathrm{kk}_{\mathcal{C}}(A,B)$ is an element represented by $\beta : J^m(A) \to B^{\mathcal{S}^m}$ then $[\alpha]\circ[\beta]$ is an element $\mathrm{kk}_{\mathcal{C}}(A,C)$ represented by

$$J^{n+m}(A) \xrightarrow{J^n(\beta)} J^n(B^{\mathcal{S}^m}) \to J^n(B)^{\mathcal{S}^m} \xrightarrow{\alpha^{\mathcal{S}^m}} C^{\mathcal{S}^{n+m}}. \tag{2.6.1}$$

This product allows us to define a composition in the category $\mathfrak{KK}_{\mathcal{C}}$ whose objects are the same objects of $\mathcal{C}$ and the morphisms from A to B are the elements of $\mathrm{kk}_{\mathcal{C}}(A,B)$. Denote by

$$j_{\mathcal{C}} : \mathcal{C} \to \mathfrak{KK}_{\mathcal{C}}$$

the functor which at the level of objects is the identity and at level of morphism sends $f : A \to B$ to $[f] \in \mathrm{kk}_{\mathcal{C}}(A,B)$.

Remark 2.6.2. A morphism $f : C \to M_\infty C$ in $\mathcal{C}$ represents an element $[f]$ in $\mathrm{kk}_{\mathcal{C}}(C, \mathcal{M}_\infty C)$. But also represents an element in $\mathrm{kk}_{\mathcal{C}}(C,C)$ because

$$\mathrm{kk}_{\mathcal{C}}(C,C) = \mathrm{colim}\, E_n(C,C)_{\mathcal{C}} \quad \text{and} \quad E_0(C,C)_{\mathcal{C}} = [C, \mathcal{M}_\infty C]_{\mathcal{C}}.$$

Consider the functor $\Omega : \mathfrak{KK}_{\mathcal{C}} \to \mathfrak{KK}_{\mathcal{C}}$ which sends an object A of $\mathcal{C}$ to the path object ΩA. Let $[\alpha]$ be an element of $\mathrm{kk}_{\mathcal{C}}(A,B)$ represented by $\alpha : J^n(A) \to B^{\mathcal{S}^n}$. The class of $\Omega[\alpha]$ is represented by

$$J^n(A^{\mathcal{S}^1}) \to J^n(A)^{\mathcal{S}^1} \xrightarrow{\alpha^{\mathcal{S}^1}} B^{\mathcal{S}^{n+1}} \tag{2.6.3}$$

Note (2.6.3) represents an element of $\mathrm{kk}_{\mathcal{C}}(A^{\mathcal{S}^1}, B^{\mathcal{S}^1})$ (see Lemma 6.3.8 [2] to check it is well defined). As $\imath : \Omega A \to A^{\mathcal{S}^1}$ is a $\mathrm{kk}_{\mathcal{C}}$-equivalence (see corollary of Lemma 6.3.2 [2]), we have the natural isomorphism

$$\mathrm{kk}_{\mathcal{C}}(A^{\mathcal{S}^1}, B^{\mathcal{S}^1}) \xrightarrow{\sim} \mathrm{kk}_{\mathcal{C}}(\Omega A, \Omega B).$$

A diagram

$$\Omega C \to A \to B \to C$$

of morphisms in $\mathfrak{KK}_{\mathcal{C}}$ is called a distinguished triangle if it is isomorphic in $\mathfrak{KK}_{\mathcal{C}}$ to the path sequence

$$\Omega B' \xrightarrow{j(\imath)} P_f \xrightarrow{j(\pi_f)} A' \xrightarrow{j(f)} B'$$

associated with a homomorphism $f : A' \to B'$ in $\mathcal{C}$. Denote the class of distinghished triangles by $\mathcal{Q}$. The category $\mathfrak{KK}_{\mathcal{C}}$ is triangulated with respect to the

endofunctor $\Omega : \mathfrak{KK}_{\mathcal{C}} \to \mathfrak{KK}_{\mathcal{C}}$ and the class $\mathcal{Q}$ of distinguished triangles, see [2, Theorem 6.5.2].

Let $\mathrm{E} : A \xrightarrow{f} B \xrightarrow{g} C$ be a weakly split extension and let $c_{\mathrm{E}} \in \mathrm{kk}_{\mathcal{C}}(J(C), A)$ be the classifying map of E. As the natural map $\rho_A : J(A) \to \Omega A$ induces a $\mathrm{kk}_{\mathcal{C}}$-equivalence (see Lemma 6.3.10, [2]) we can consider the following morphisms in $\mathrm{kk}_{\mathcal{C}}(\Omega C, A)$

$$\partial_{\mathrm{E}} := c_{\mathrm{E}} \circ \rho_C^{-1}. \tag{2.6.4}$$

The functor $j_{\mathcal{C}} : \mathcal{C} \to \mathfrak{KK}_{\mathcal{C}}$ with the morphisms $\{\partial_{\mathrm{E}} : \mathrm{E} \in \mathcal{E}\}$ is an excisive homology theory, homotopy invariant and M_∞-stable.

Theorem 2.6.5. *The functor $j_{\mathcal{C}} : \mathcal{C} \to \mathfrak{KK}_{\mathcal{C}}$ is universal with the properties defined above. In other words, if $\mathcal{T}$ is a triangulated category and $G : \mathcal{C} \to \mathcal{T}$ together a class of morphisms $\{\overline{\partial}_{\mathrm{E}} : \mathrm{E} \in \mathcal{E}\}$ is an excisive, homotopy invariant and M_∞-stable functor, then there exists a unique triangle functor $\overline{G} : \mathfrak{KK}_{\mathcal{C}} \to \mathcal{T}$ such that the following diagram commutes*

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{j_{\mathcal{C}}} & \mathfrak{KK}_{\mathcal{C}} \\ & \searrow^{G} & \downarrow{\scriptstyle \overline{G}} \\ & & \mathcal{T} \end{array}$$

Proof. By definition $j_{\mathcal{C}}$ is homotopy invariant and M_∞-stable. Let us show that $j_{\mathcal{C}} : \mathcal{C} \to \mathfrak{KK}_{\mathcal{C}}$ is an excisive homology theory with the family $\{\partial_{\mathrm{E}} : \mathrm{E} \in \mathcal{E}\}$. Let $\mathrm{E} : A \xrightarrow{f} B \xrightarrow{g} C$ be a weakly split extension. Take the path sequence asociated to g and the following diagram in $\mathfrak{KK}_{\mathcal{C}}$

$$\begin{array}{ccccccccc} \Omega C & \xrightarrow{j(\iota)} & P_g & \xrightarrow{j(\pi_g)} & B & \xrightarrow{j(g)} & C & & T \\ \uparrow{\scriptstyle \mathrm{id}} & & \uparrow{\scriptstyle \iota_f} & & \uparrow{\scriptstyle \mathrm{id}} & & \uparrow{\scriptstyle \mathrm{id}} & & \\ \Omega C & \xrightarrow[\partial_{\mathrm{E}}]{} & A & \xrightarrow[j(f)]{} & B & \xrightarrow[j(g)]{} & C & & T' \end{array}$$

The first square commutes beacuse $\iota_f \circ c_{\mathrm{E}}$ is elementarily homotopic to $\iota \circ \rho_C$. By [2, Lemma 6.3.2], the morphism ι_f is a $\mathrm{kk}_{\mathcal{C}}$-equivalence. Finally T and T' are isomorphic in $\mathfrak{KK}_{\mathcal{C}}$ and T' is a distinguished triangle. For the rest of the proof see [2, Theorem 6.6.2]. □

Algebraic K-theory is not excisive, nor is it M_∞-stable or homotopy invariant. However the homotopy algebraic K-theory, defined in [12] and denoted by KH, have all these properties. Consider the funtor $\mathrm{KH} = \mathrm{KH}_0 : \mathrm{Alg}_\ell \to \mathrm{Ab}$. Because it satisfies [2, Theorem 6.6.6] we have a natural map

$$\mathrm{kk}_{\mathrm{Alg}_\ell}(\ell, A) \to \hom_{\mathrm{Ab}}(\mathrm{KH}(\ell), \mathrm{KH}(A))$$

As ℓ is unital, there is a map $\mathbb{Z} \to \ell$ which induce a map

$$\hom_{\mathrm{Ab}}(\mathrm{KH}(\ell), \mathrm{KH}(A)) \mapsto \hom_{\mathrm{Ab}}(\mathrm{KH}(\mathbb{Z}), \mathrm{KH}(A)) \simeq \mathrm{KH}(A).$$

Composing both maps we obtain a homomorphism $\mathrm{kk}_{\mathrm{Alg}_\ell}(\ell, A) \to \mathrm{KH}(A)$ and the main theorem from [2] prove that it is an isomorphism.

Theorem 2.6.6. *Consider $\mathcal{C} = \mathrm{Alg}_\ell$ the category of ℓ-algebras, then*

$$\mathrm{kk}_{\mathrm{Alg}_\ell}(\ell, A) \simeq \mathrm{KH}(A).$$

Proof. See [2, Theorem 8.2.1]. □

3. Equivariant matrix invariance

In the equivariant setting we replace the property of M_∞-stability by a stability condition depending on $\mathcal{C}$. We consider the different cases of $\mathcal{C}$ separately.

3.1. G-equivariant stability.

Regard

$$M_G = \{f : G \times G \to \ell : \mathrm{supp}(f) < \infty\}$$

as the algebra of matrices with coefficients in ℓ indexed by $G \times G$, with translation action of G:

$$g \cdot e_{s,t} = e_{gs,gt}.$$

We are going to identify a G-algebra A with the G-algebra $M_G \otimes A$ with the diagonal action. Note that the map $a \to a \otimes e_{1_G,1_G}$ is not equivariant then we can not define G-stability as in Section 2.2. For this reason we define G-stability in terms of G-modules.

A pair $(\mathcal{W}, B)$ is a G-module with basis if $\mathcal{W}$ is a G-module, free as an ℓ-module and B is a basis of $\mathcal{W}$. A pair $(\mathcal{W}', B')$ is a submodule with basis of $(\mathcal{W}, B)$ if $\mathcal{W}'$ is a submodule of $\mathcal{W}$ and $B' \subset B$. Note that if $(\mathcal{W}_1, B_1)$ and $(\mathcal{W}_2, B_2)$ are G-modules with basis then $(\mathcal{W}_1 \oplus \mathcal{W}_2, B_1 \sqcup B_2)$ is a G-module with basis.

Let $(\mathcal{W}, B)$ be a G-module with basis B. We define

$$\mathcal{L}(\mathcal{W}, B) := \{\psi : B \times B \to \ell : \{v : \psi(v, w) \neq 0\} \text{ is finite for all } w\}$$

Note that $\mathcal{L}(\mathcal{W}, B)$ and $\mathrm{End}_\ell(\mathcal{W}) = \{f : \mathcal{W} \to \mathcal{W} : \ f \text{ is } \ell\text{-linear}\}$ are isomorphic; indeed we have inverse isomorphisms

$$\mathrm{End}_\ell(\mathcal{W}) \to \mathcal{L}(\mathcal{W}, B) \qquad f \mapsto \psi_f \qquad \psi_f(v, w) = p_v(f(w)) \tag{3.1.1}$$

here $p_v : \mathcal{W} \to \ell$ is the projection to the submodule of $\mathcal{W}$ generated by v

$$\mathcal{L}(\mathcal{W}, B) \to \mathrm{End}_\ell(\mathcal{W}) \qquad \psi \mapsto f_\psi \qquad f_\psi(w) = \sum_{v \in B} \psi(v, w)v. \tag{3.1.2}$$

Define

$$\begin{aligned}
&\mathcal{C}(\mathcal{W}, B) := \{\psi \in \mathcal{L}(\mathcal{W}, B) : \{w : \psi(v, w) \neq 0\} \text{ is finite for all } v\}\\
&\mathcal{F}(\mathcal{W}, B) := \{\psi \in \mathcal{L}(\mathcal{W}, B) : \{(v, w) : \psi(v, w) \neq 0\} \text{ is finite } \}\\
&\mathrm{End}^F_\ell(\mathcal{W}, B) := \{f \in \mathrm{End}_\ell(\mathcal{W}) : \psi_f \in \mathcal{F}(\mathcal{W}, B)\}\\
&\mathrm{End}^C_\ell(\mathcal{W}, B) := \{f \in \mathrm{End}_\ell(\mathcal{W}) : \psi_f \in \mathcal{C}(\mathcal{W}, B)\}
\end{aligned}$$

Note that $\mathcal{C}(\mathcal{W}, B)$ is a ring with the matrix product and $\mathrm{End}^C_\ell(\mathcal{W}, B)$ is a ring with the composition. These rings are isomorphic.

Let $(\mathcal{W}, B)$ be a G-module with basis. Consider the representation

$$\rho : G \to \mathrm{End}_\ell(\mathcal{W}) \qquad\qquad \rho_g(w) = g \cdot w$$

We say that $(\mathcal{W}, B)$ is a G-module by finite automorphisms if $\rho(G) \subset \mathrm{End}^F_\ell(\mathcal{W}, B)$. We say that $(\mathcal{W}, B)$ is a G-module by locally finite automorphisms if $\rho(G) \subset \mathrm{End}^C_\ell(\mathcal{W}, B)$. If $(\mathcal{W}, B)$ is a G-module by locally finite automorphisms, $\mathrm{End}^C_\ell(\mathcal{W}, B)$ and $\mathrm{End}^F_\ell(\mathcal{W}, B)$ are G-algebras with the following action

$$g \cdot f = \rho(g) f(\rho(g))^{-1}$$

Note that $\mathrm{End}^F_\ell(\mathcal{W}, B)$ is an ideal of $\mathrm{End}^C_\ell(\mathcal{W}, B)$.

Example 3.1.3. Let $\mathcal{W} = \ell G$ be the group algebra considered as a G-module via the regular representation with basis $B = \{\delta_g : g \in G\}$,

$$g \cdot (\sum_{h\in G} a_h \delta_h) = \sum_{h\in G} a_h \delta_{gh}, \qquad a_h \in \ell. \tag{3.1.4}$$

Note

$$\rho : G \to \mathrm{End}_\ell(\ell G) \simeq \mathcal{L}(\ell G, B) \quad g \mapsto M_g = \sum_{t\in G} e_{gt,t}$$

As $M_g \in \mathcal{C}(\ell G, B)$ for all $g \in G$, $(\ell G, B)$ is a G-module by locally finite automorphisms. Moreover we have $(M_g)^{-1} = M_{g^{-1}} = (M_g)^t$ and $M_G = \mathcal{F}(\ell G, B)$.

Let A be a G-algebra. Consider the tensor product $M_G A = M_G \otimes A$ with the diagonal action of G.

Remark 3.1.5. Let $\mathcal{W}$ be a G-module. Let $\mathcal{W}^\tau$ be $\mathcal{W}$ considered as a G-module with trivial action. Recall that $\ell G \otimes \mathcal{W} \simeq \ell G \otimes \mathcal{W}^\tau$. Inverse isomorphisms are given by

$$T : \ell G \otimes \mathcal{W}^\tau \to \ell G \otimes \mathcal{W} \qquad S : \ell G \otimes \mathcal{W} \to \ell G \otimes \mathcal{W}^\tau$$

$$T(\delta_g \otimes h) = \delta_g \otimes g(h) \qquad S(\delta_g \otimes h) = \delta_g \otimes g^{-1}(h)$$

If $(\mathcal{W}, B)$ is a G-module with basis we will write $\mathrm{End}_\ell^C(\mathcal{W})$ and $\mathrm{End}_\ell^F(\mathcal{W})$ ommiting the basis when there is no confusion.

Remark 3.1.6. Let $(\mathcal{W}, B)$ be a G-module with basis. Let us check that

$$\mathrm{End}_\ell^F(\ell G \otimes \mathcal{W}) \simeq \mathrm{End}_\ell^F(\ell G) \otimes \mathrm{End}_\ell^F(\mathcal{W}).$$

Define a G-algebra homomorphism $T : \mathrm{End}_\ell^F(\ell G) \otimes \mathrm{End}_\ell^F(\mathcal{W}) \to \mathrm{End}_\ell^F(\ell G \otimes \mathcal{W})$ by

$$T(e_{g,h} \otimes e_{v,w}) = e_{g,v,h,w} \qquad v, w \in B \quad g, h \in G$$

As T is a bijection between the basis, T is an isomorphism.

Let $(\mathcal{W}_1, B_1)$ and $(\mathcal{W}_2, B_2)$ be G-modules by locally finite automorphisms such that $\mathrm{card}(B_i) \leq \mathrm{card}(G) \times \mathrm{card}\,\mathbb{N}$, $i = 1, 2$. The inclusion $\iota : \mathcal{W}_1 \to \mathcal{W}_1 \oplus \mathcal{W}_2$ induces a morphism of G-algebras

$$\tilde{\iota} : \mathrm{End}_\ell^F(\mathcal{W}_1) \to \mathrm{End}_\ell^F(\mathcal{W}_1 \oplus \mathcal{W}_2) \qquad f \mapsto \begin{pmatrix} f & 0 \\ 0 & 0 \end{pmatrix} \tag{3.1.7}$$

Let A be a G-algebra and consider

$$\tilde{\iota} \otimes 1 : \mathrm{End}_\ell^F(\mathcal{W}_1) \otimes A \to \mathrm{End}_\ell^F(\mathcal{W}_1 \oplus \mathcal{W}_2) \otimes A.$$

A functor $F : G\text{-Alg} \to \mathcal{D}$ is G-stable if for $(\mathcal{W}_1, B_1)$, $(\mathcal{W}_2, B_2)$ and A as above $F(\tilde{\iota} \otimes 1)$ is an isomorphism in $\mathcal{D}$.

Let us show that if $F : G\text{-Alg} \to \mathcal{D}$ is a G-stable functor then F is M_∞-stable. Consider $(\mathcal{W}_1, B_1) = (\ell, \{1\})$ and $(\mathcal{W}_2, B_2) = (\ell^{(\mathbb{N})}, \{e_i : i \in \mathbb{N}\})$ with $\ell^{(\mathbb{N})} = \oplus_{i=1}^\infty \ell$, $\{e_i : i \in \mathbb{N}\}$ is the canonical basis and both modules have the trivial action of G. Note

$$\mathrm{End}_\ell^F(\ell) = \mathrm{End}_\ell(\ell) = \ell \quad \mathrm{End}_\ell^F(\ell \oplus \ell^{(\mathbb{N})}) = \mathrm{End}_\ell^F(\ell^{(\mathbb{N})}) = M_\infty$$

and $\tilde{\iota}_\infty : \ell \to M_\infty$ is the inclusion at the upper left corner. Then $\tilde{\iota} \otimes 1 = \iota : A \to M_\infty(A)$ and $F(\iota)$ is an isomorphism. Observe that if $G = \{e\}$, F is G-stable if and only if F is M_∞-stable.

Let A, B be G-algebras and $F : G\text{-Alg} \to \mathcal{D}$ a functor. A zig-zag between A and B by F is a diagram in G-Alg

$$A \xrightarrow{f_1} C_1 \xleftarrow{g_1} \dots \xrightarrow{f_n} C_n \xleftarrow{g_n} B$$

such that $F(g_i)$, $i = 1, \dots, n$, are isomorphisms in $\mathcal{D}$.

Example 3.1.8. Let A be a G-algebra and F a G-stable functor. There exists a zig-zag between A and $M_G A$ by F. Consider $\mathcal{W}_1 = (\ell G, B)$ as in the example 3.1.3 and consider $\mathcal{W}_2 = (\ell, \{1\})$ with the trivial action of G. Put $\mathcal{W} = \mathcal{W}_1 \oplus \mathcal{W}_2$ and $C = \mathrm{End}_\ell^F(\mathcal{W})$ with the induced action, then

$$\iota : A = A \otimes \ell = A \otimes \mathrm{End}_\ell^F(\ell) \to A \otimes C \leftarrow A \otimes M_G : \iota'$$

is a zig-zag between A and $M_G A$ by F.

Proposition 3.1.9. *Suppose G is countable. Let $F : G\text{-Alg} \to \mathcal{D}$ be an M_∞-stable functor. Define*

$$\hat{F} : G\text{-Alg} \to \mathcal{D} \qquad A \mapsto F(M_G \otimes A)$$

Then $\hat{F}$ is G-stable.

Proof. Let $(\mathcal{W}_1, B_1)$, $(\mathcal{W}_2, B_2)$ be G-modules by locally finite automorphisms and let A be a G-algebra. Consider

$$\tilde{\iota} \otimes 1 : \mathrm{End}_\ell^F(\mathcal{W}_1) \otimes A \to \mathrm{End}_\ell^F(\mathcal{W}_1 \oplus \mathcal{W}_2) \otimes A.$$

We have to prove that

$$\hat{F}(\tilde{\iota} \otimes 1) : F(M_G \otimes \mathrm{End}_\ell^F(\mathcal{W}_1) \otimes A) \to F(M_G \otimes \mathrm{End}_\ell^F(\mathcal{W}_1 \oplus \mathcal{W}_2) \otimes A) \tag{3.1.10}$$

is an isomorphism. By remarks 3.1.6 and 3.1.5 we know that

$$F(M_G \otimes \mathrm{End}_\ell^F(\mathcal{W}_1) \otimes A) \simeq F(\mathrm{End}_\ell^F(\mathcal{W}_1^\tau) \otimes M_G \otimes A)$$

and

$$F(M_G \otimes \mathrm{End}_\ell^F(\mathcal{W}_1 \oplus \mathcal{W}_2) \otimes A) \simeq F(\mathrm{End}_\ell^F((\mathcal{W}_1 \oplus \mathcal{W}_2)^\tau) \otimes M_G \otimes A).$$

Note that $\mathrm{End}_\ell^F(\mathcal{W}_1^\tau)$ and $\mathrm{End}_\ell^F((\mathcal{W}_1 \oplus \mathcal{W}_2)^\tau)$ are equivariantly isomorphic to M_n or M_∞. As F is M_∞-stable, (3.1.10) is an isomorphism. □

Remark 3.1.11. Suppose G is a finite group of order n such that $1/n \in \ell$. Then the element $\xi = (1/n)\sum_{g \in G} \delta_g$ in ℓG is idempotent. The map $s : \ell \to \ell G$, $s(1) = \xi$, is a G-equivariant section of the canonical augmentation $\varphi : \ell G \to \ell$. Thus the sequence of G-modules

$$0 \longrightarrow I \longrightarrow \ell G \xrightarrow{\varphi} \ell \longrightarrow 0 \tag{3.1.12}$$

splits. Hence $\ell G = \ell\xi \oplus I$. Notice that I is a G-module with basis $\{\delta_e - \delta_g : g \neq e\}$. Define

$$\lambda_g = \begin{cases} \xi & g = e \\ \delta_e - \delta_g & g \neq e \end{cases}$$

The set $\Lambda = \{\lambda_g : g \in G\}$ is a basis of ℓG and the relations with the elements of $B = \{\delta_g : g \in G\}$ are the following

$$\begin{array}{lll} \lambda_e = \frac{1}{n}\sum_{g \in G} \delta_g & \lambda_h = \delta_e - \delta_h & \\ \delta_e = \lambda_e + \frac{1}{n}\sum_{g \neq e} \lambda_g & \delta_h = \lambda_e - \lambda_h + \frac{1}{n}\sum_{g \neq e} \lambda_g & h \neq e \end{array}$$

Consider $\mathcal{W}_1 = \ell = (\ell\xi, \{\xi\})$ and $\mathcal{W}_2 = (I, \{\lambda_g\}_{g\neq e})$, in this case the morphism (3.1.7) is

$$\iota : \ell \to M_G \simeq \mathrm{End}(\ell G, \Lambda) \quad 1 \mapsto \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 0 \end{pmatrix}$$

If we write it in the canonical basis B we have

$$\overline{\iota} : \ell \to M_G = \mathrm{End}(\ell G, B) \quad 1 \mapsto \begin{pmatrix} \frac{1}{n} & \frac{1}{n} & \dots & \frac{1}{n} \\ \frac{1}{n} & \frac{1}{n} & \dots & \frac{1}{n} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{1}{n} & \frac{1}{n} & \dots & \frac{1}{n} \end{pmatrix}$$

If $F : G\text{-Alg} \to \mathcal{D}$ is a G-stable functor then $F(\overline{\iota})$ is an isomorphism in $\mathcal{D}$.

3.2. **G-graded stability.** In this section we consider a dual notion of G-equivariant stability. We want to identify a G-graded algebra A with the G-graded matrix algebra $M_G A$. The definition of G-graded stability is easier than that of G-equivariant stability because the morphism $A \to M_G A$, $a \mapsto e_{1_G.1_G} \otimes a$, is homogeneous.

Write G_{gr}-Alg for the category of G-graded algebras. Let A be a G-graded algebra. Define the following grading in $M_G A$

$$(M_G A)_g := < e_{s,t} \otimes a : g = s|a|t^{-1} > . \tag{3.2.1}$$

Depending on the context $M_G A$ will be considered as an algebra, a G-algebra or a G-graded algebra. Let be

$$\iota_A : A \to M_G A \qquad a \mapsto e_{1_G,1_G} \otimes a,$$

note ι_A is homogeneous. A functor $F : G_{gr}\text{-Alg} \to \mathcal{D}$ is G_{gr}-stable if $F(\iota_A)$ is an isomorphism in $\mathcal{D}$ for all $A \in G_{gr}$-Alg.

Proposition 3.2.2. *Suppose G is countable. Let $F : G_{gr}\text{-Alg} \to \mathcal{D}$ be an M_∞-stable functor. Define*


$$\hat{F} : G_{gr}\text{-Alg} \to \mathcal{D} \qquad A \mapsto F(M_G A)$$

with $M_G A$ as in (3.2.1)*. Then $\hat{F}$ is G_{gr}-stable.*

Proof. Denote by $M_{|G|}$ to M_G with the trivial grading. Because G is countable we have that $M_\infty M_{|G|}$ and M_∞ are isomorphic in G_{gr}-Alg. We also have the following isomorphisms of G-graded algebras

$$\begin{aligned} \eta : M_G M_G \to M_G M_{|G|} \qquad & e_{s,g} \otimes e_{t,r} \mapsto e_{st,gr} \otimes e_{t,r} \\ \mu : M_G M_{|G|} \to M_{|G|} M_G \qquad & e_{s,g} \otimes e_{t,r} \mapsto e_{t,r} \otimes e_{s,g} \end{aligned}$$

Consider the following commutative diagram



It follows that $\hat{F}(\iota_A) = F(\mathrm{id}_{M_G} \otimes \iota_A)$ is an isomorphism in $\mathcal{D}$ because F is M_∞-stable. □

4. Equivariant algebraic kk-theory

From now to the rest of the paper we suppose G is a countable group.

4.1. **The category $\mathfrak{KK}^G$.** In this section we introduce an equivariant algebraic kk-theory for G-algebras. Let A, B be G-algebras, we define

$$\mathrm{kk}^G(A,B) := \mathrm{kk}_{G\text{-Alg}}(M_G \otimes A, M_G \otimes B).$$

Consider the category $\mathfrak{KK}^G$ whose objects are the G-algebras and where the morphisms between A and B are the elements of $\mathrm{kk}^G(A,B)$. Let $j^G : G\text{-Alg} \to \mathfrak{KK}^G$ be the functor defined as the identity on objects and which sends each morphism of G-algebras $f : A \to B$ to its class $[\mathrm{id}_{M_G} \otimes f] \in \mathrm{kk}^G(A,B)$. The composition law in $\mathfrak{KK}^G$ is the same that in (2.6.1) taking $M_G \otimes A$, $M_G \otimes B$ and $M_G \otimes C$ instead of A, B and C. As in Section 2.6 we have an equivalence $\Omega : \mathfrak{KK}^G \to \mathfrak{KK}^G$ and distinguished triangles

$$\Omega C \to A \to B \to C$$

in $\mathfrak{KK}^G$ which gives to $\mathfrak{KK}^G$ a triangulated category structure.

Theorem 4.1.1. *The functor $j^G : G\text{-Alg} \to \mathfrak{KK}^G$ is an excisive, equivariantly homotopy invariant, and G-stable functor. Moreover, it is the universal functor for these properties. In other words, if $\mathcal{T}$ is a triangulated category and $R : G\text{-Alg} \to \mathcal{T}$ together a class of morphisms $\{\overline{\partial}_{\mathrm{E}} : \mathrm{E} \in \mathcal{E}\}$ is an excisive, equivariantly homotopy invariant and G-stable functor, then there exists a unique triangle functor $\overline{R} : \mathfrak{KK}^G \to \mathcal{T}$ such that the following diagram commutes*

$$\begin{array}{ccc} G\text{-Alg} & \xrightarrow{j^G} & \mathfrak{KK}^G \\ & \searrow^{R} & \downarrow{\scriptstyle \overline{R}} \\ & & \mathcal{T} \end{array}$$

Proof. Let E be a weakly split extension. Define


$$\partial^G_{\mathrm{E}} \in \hom_{\mathfrak{KK}^G}(\Omega C, A) = \hom_{\mathfrak{KK}_{G\text{-Alg}}}(\Omega M_G \otimes C, M_G \otimes A)$$

as the morphism $\partial_{\mathrm{E}'}$ defined in (2.6.4) asociated to the following weakly split extension

$$M_G \otimes A \to M_G \otimes B \to M_G \otimes C \qquad (\mathrm{E}')$$

By theorem 2.6.5 and proposition 3.1.9 the functor $j^G : G\text{-Alg} \to \mathfrak{KK}^G$ with the family $\{\partial^G_{\mathrm{E}} : \mathrm{E} \in \mathcal{E}\}$ is an excisive, homotopy invariant and G-stable functor. Let us check it is universal for these properties. Let $X : G\text{-Alg} \to \mathcal{T}$ be a functor which has the mentioned properties with a family $\{\overline{\partial}_{\mathrm{E}} : \mathrm{E} \in \mathcal{E}\}$. By theorem 2.6.5 there exists a unique triangle functor $\overline{X} : \mathfrak{KK}_{G\text{-Alg}} \to \mathcal{T}$ such that the following diagram

commutes

(4.1.2)


We will define $X' : \mathfrak{KK}^G \to \mathcal{T}$. We know that $X' = X$ on objects. As X is G-stable the following morphisms are a zig-zag between A and $M_G \otimes A$ by X, (see Example 3.1.8)

$$A \xrightarrow{\iota_A} M_{G\sqcup\{*\}} \otimes A \xleftarrow{\iota'_A} M_G \otimes A \tag{4.1.3}$$

Let $\alpha \in \mathrm{kk}^G(A, B)$ and define

$$X'(\alpha) := X(\iota_B)^{-1} X(\iota'_B) \overline{X}(\alpha) X(\iota'_A)^{-1} X(\iota_A).$$


Note this definition is the unique possibility to make the diagram (4.1.2) commutative. □

4.2. **The category $\hat{\mathfrak{KK}}^G$.** In this section we introduce the equivariant algebraic kk-theory for G-graded algebras. Let A, B be G-graded algebras. We define

$$\hat{\mathrm{kk}}^G(A, B) := \mathrm{kk}_{G_{gr}\text{-Alg}}(M_G A, M_G B).$$

Consider the category $\hat{\mathfrak{KK}}^G$ whose objects are the G-graded algebras and the morphisms between A and B are the elements of $\hat{\mathrm{kk}}^G(A, B)$. Let $j^{G_{gr}} : G_{gr}\text{-Alg} \to \hat{\mathfrak{KK}}^G$ be the functor defined as the identity on objects and which sends each morphism of G-graded algebras $f : A \to B$ to its class $[\mathrm{id}_{M_G} f] \in \hat{\mathrm{kk}}^G(A, B)$. As in Section 4.1 we can consider a triangulated category structure on $\hat{\mathfrak{KK}}^G$.

Theorem 4.2.1. *The functor $j^{G_{gr}} : G_{gr}\text{-Alg} \to \hat{\mathfrak{KK}}^G$ is an excisive, graded homotopy invariant, and G-graded stable functor. Moreover, it is the universal functor for these properties. In other words, if $\mathcal{T}$ is a triangulated category and $R : G_{gr}\text{-Alg} \to \mathcal{T}$ together a class of morphisms $\{\overline{\partial}_{\mathrm{E}} : \mathrm{E} \in \mathcal{E}\}$ is an excisive, graded homotopy invariant and G-graded stable functor, then there exists a unique triangle functor $\overline{R} : \hat{\mathfrak{KK}}^G \to \mathcal{T}$ such that the following diagram commutes*



Proof. The proof is similar to Theorem 4.1.1. □

5. Algebraic Green-Julg theorem

5.1. **Crossed product and trivial action.** Let A be an ℓ-algebra. Write A^τ for A with the trivial action of G. This gives us a functor $\tau : \mathrm{Alg} \to G\text{-Alg}$. It is easy to check that $j^G \circ \tau$ satisfies excision, is homotopy invariant and is M_∞-stable. Write $\mathfrak{KK} = \mathfrak{KK}_{\mathrm{Alg}}$. By Theorem 2.6.5 there exists a unique functor $\tau : \mathfrak{KK} \to \mathfrak{KK}^G$ such that the following diagram is commutative

$$\begin{array}{ccc} \mathrm{Alg} & \xrightarrow{\tau} & G\text{-Alg} \\ {\scriptstyle j}\downarrow & & \downarrow{\scriptstyle j^G} \\ \mathfrak{KK} & \xrightarrow[\tau]{} & \mathfrak{KK}^G \end{array}$$

Let A be a G-algebra. The crossed product algebra $A \rtimes G$ is the ℓ-module $A \otimes \ell G$ with the following multiplication

$$(a \rtimes g)(b \rtimes h) = a(g \cdot b) \rtimes gh \qquad a, b \in A \ g, h \in G.$$

Proposition 5.1.1. *Let A be a G-algebra and $\mathcal{W}$ a G-module by locally finite automorphisms. The following algebras are naturally isomorphic*

$$(A \rtimes G) \otimes \mathrm{End}^F_\ell(\mathcal{W}) \simeq (A \otimes \mathrm{End}^F_\ell(\mathcal{W})) \rtimes G.$$

Proof. Let $\rho : G \to (\mathrm{End}^K_\ell(\mathcal{W}))^\times$ be the structure map. Note that the homomorphisms

$$\phi : (A \rtimes G) \otimes \mathrm{End}^F_\ell(\mathcal{W}) \to (A \otimes \mathrm{End}^F_\ell(\mathcal{W})) \rtimes G \qquad \phi(a \rtimes g \otimes \varphi) = a \otimes \varphi\rho(g^{-1}) \rtimes g$$

$$\psi : (A \otimes \mathrm{End}^F_\ell(\mathcal{W})) \rtimes G \to (A \rtimes G) \otimes \mathrm{End}^F_\ell(\mathcal{W}) \qquad \psi(a \otimes \varphi \rtimes g) = a \rtimes g \otimes \varphi\rho(g)$$

are inverse of each other. □

Proposition 5.1.2. *There exists a unique functor $\rtimes G : \mathfrak{KK}^G \to \mathfrak{KK}$ such that the following diagram is commutative*

$$\begin{array}{ccc} G\text{-Alg} & \xrightarrow{\rtimes G} & \mathrm{Alg} \\ {\scriptstyle j^G}\downarrow & & \downarrow{\scriptstyle j} \\ \mathfrak{KK}^G & \xrightarrow[\rtimes G]{} & \mathfrak{KK} \end{array}$$

Proof. We shall show $j(- \rtimes G)$ is excisive, homotopy invariant and G-stable. Because $\rtimes G$ maps split exact sequences to split exact sequences and j is excisive, then $j(- \rtimes G)$ is excisive.

That $j(- \rtimes G)$ is homotopy invariant follows from the fact that

$$A[t] \rtimes G = (A \rtimes G)[t].$$

Let $(\mathcal{W}_1, B_1)$, $(\mathcal{W}_2, B_2)$ be G-modules by locally finite automorphisms and A a G-algebra. Consider the isomorphism ψ defined in Proposition 5.1.1. Note that the following diagram is commutative

$$\begin{array}{ccc} (A \otimes \mathrm{End}^F_\ell(\mathcal{W}_1)) \rtimes G & \xrightarrow{(1 \otimes \tilde{\iota}) \rtimes G} & (A \otimes \mathrm{End}^F_\ell(\mathcal{W}_1 \oplus \mathcal{W}_2)) \rtimes G \\ {\scriptstyle \psi}\downarrow & & \downarrow{\scriptstyle \psi} \\ (A \rtimes G) \otimes \mathrm{End}^F_\ell(\mathcal{W}_1) & \xrightarrow[(1 \rtimes G) \otimes \tilde{\iota}]{} & (A \rtimes G) \otimes \mathrm{End}^F_\ell(\mathcal{W}_1 \oplus \mathcal{W}_2) \end{array}$$

Because j is M_∞-stable, $j((1 \rtimes G) \otimes \tilde{\imath})$ is an isomorphism. Hence $j(- \rtimes G)(1 \otimes \tilde{\imath})$ is an isomorphism by the diagram above. □

Remark 5.1.3. Let $[\iota_\infty \alpha] \in \mathrm{kk}^G(A, B)$ be an element represented by $\alpha : J^n(M_G A) \to (M_G B)^{\mathrm{sd}^p \mathcal{S}^n}$ which is a morphism in $[J^n(M_G A), \mathcal{M}_\infty(M_G B)^{\mathrm{sd}^p \mathcal{S}^n}]$. Consider the classifying map

$$J^n(M_G A \rtimes G) \to J^n(M_G A) \rtimes G$$

The element $[\alpha] \rtimes G$ is represented by the following composition

$$J^n(M_G A \rtimes G) \to J^n(M_G A) \rtimes G \xrightarrow{\alpha \rtimes G} (M_G B)^{\mathrm{sd}^p \mathcal{S}^n} \rtimes G.$$

Proposition 5.1.4. *The functor $\rtimes G : \mathfrak{KK}^G \to \mathfrak{KK}$ is a triangle functor.*

Proof. A distinguished triangle in $\mathfrak{KK}^G$ is a diagram isomorphic to

$$\Omega B \xrightarrow{j^G(\iota)} P_f \xrightarrow{j^G(\pi_f)} A \xrightarrow{j^G(f)} B$$

for some morphism of G-algebras $f : A \to B$. That means, if it is isomophic to

$$M_G \Omega B \xrightarrow{j_{G\text{-Alg}}(\iota)} M_G P_f \xrightarrow{j_{G\text{-Alg}}(\pi_f)} M_G A \xrightarrow{j_{G\text{-Alg}}(f)} M_G B \tag{5.1.5}$$

in $\mathfrak{KK}_{G\text{-Alg}}$. The functor $\rtimes G : \mathfrak{KK}^G \to \mathfrak{KK}$ sends the triangle (5.1.5) to

$$(M_G \Omega B) \rtimes G \to (M_G P_f) \rtimes G \to (M_G A) \rtimes G \to (M_G B) \rtimes G \tag{5.1.6}$$

which by Proposition 5.1.1 and M_∞-stability is isomorphic to

$$\Omega(B \rtimes G) \to P_f \rtimes G \to A \rtimes G \to B \rtimes G \tag{5.1.7}$$

As $P_f \rtimes G \simeq P_{f \rtimes G}$, (5.1.7) is a distinghished triangle in $\mathfrak{KK}$. □

5.2. **Green-Julg Theorem for $\mathfrak{KK}^G$.** In this section we shall see an algebraic version of the Green-Julg Theorem, see [11] and [6] for versions of this result in Kasparov KK-theory and E-theory setting.

Theorem 5.2.1. *Let G be a finite group of n elements and $1/n \in \ell$. The functors $\tau : \mathfrak{KK} \to \mathfrak{KK}^G$ and $\rtimes G : \mathfrak{KK}^G \to \mathfrak{KK}$ are adjoint functors. Hence*

$$\mathrm{kk}^G(A^\tau, B) \simeq \mathrm{kk}(A, B \rtimes G) \qquad A \in \mathrm{Alg} \qquad B \in G\text{-Alg}.$$

Proof. By [10, Theorem 2, pag 81], it is enough to prove that there exist natural transformations $\overline{\alpha}_A \in \mathrm{kk}(A, A^\tau \rtimes G)$ and $\overline{\beta}_B \in \mathrm{kk}^G((B \rtimes G)^\tau, B)$ such that the following compositions

$$A^\tau \xrightarrow{\tau(\overline{\alpha}_A)} (A^\tau \rtimes G)^\tau \xrightarrow{\overline{\beta}_{\tau(A)}} A^\tau \qquad B \rtimes G \xrightarrow{\overline{\alpha}_{B \rtimes G}} (B \rtimes G)^\tau \rtimes G \xrightarrow{\overline{\beta}_B \rtimes G} B \rtimes G$$

are the identities in $\mathrm{kk}^G(A^\tau, A^\tau)$ and $\mathrm{kk}(B \rtimes G, B \rtimes G)$ respectively.

Put $\epsilon = 1/n \sum_{g \in G} g$ in ℓG and define

$$\alpha_A : A \to A^\tau \rtimes G = A \otimes \ell G \qquad \alpha(a) = a \otimes \epsilon. \tag{5.2.2}$$

Note α_A is an algebra morphism since ϵ is idempotent. Consider the element $\overline{\alpha}_A \in \mathrm{kk}(A, A^\tau \rtimes G)$ represented by α_A. Let

$$\beta_B : (B \rtimes G)^\tau \to M_G B \qquad \beta_B(b \rtimes g) = \sum_{s \in G} s(b) e_{s, sg}$$

One can check that β_B is an equivariant algebra morphism. Let $\overline{\beta}_B \in \mathrm{kk}^G((B \rtimes G)^\tau, B)$ be represented by β_B. The composite $\beta_{\tau(A)} \tau(\alpha_A)$ is $\mathrm{id}_{A^\tau} \otimes \overline{\iota}$ where $\overline{\iota}$ is the map defined in 3.1.11. As j^G is G-stable, $j^G(\mathrm{id}_{A^\tau} \otimes \overline{\iota})$ is the identity in $\mathrm{kk}^G(A^\tau, A^\tau)$.

Let ψ be the morphism defined in the proof of the Proposition 5.1.1. By Remark 5.1.3, $\overline{\beta}_B \rtimes G$ is represented by $\psi \circ (\beta_B \rtimes G)$. We want to prove that $\psi \circ (\beta_B \rtimes G) \circ \alpha_{B\rtimes G}$ represents the identity in $\mathrm{kk}(B \rtimes G, B \rtimes G)$. Note that

$$(\psi \circ (\beta_B \rtimes G) \circ \alpha_{B\rtimes G})(b \rtimes g) = \frac{1}{n} \sum_{h,s\in G} (s(b) \rtimes h) e_{s,h^{-1}sg}.$$

Put $t = h^{-1}sg$ and note

$$\frac{1}{n} \sum_{h,s\in G} (s(b) \rtimes h) e_{s,h^{-1}sg} = \frac{1}{n} \sum_{t,s\in G} (s(b) \rtimes sgt^{-1}) e_{s,t} \tag{5.2.3}$$

and

$$s(b) \rtimes sgt^{-1} = (\mathbf{1} \rtimes s)(b \rtimes g)(\mathbf{1} \rtimes t^{-1}) \qquad \text{in } \tilde{B} \rtimes G$$

We can write (5.2.3) as $TA_{b\rtimes g}T^{-1}$, where

$$A_{b\rtimes g} = \frac{1}{n} \sum_{t,s\in G} (b \rtimes g) e_{s,t} \quad T = \sum_{t\in G} (\mathbf{1} \rtimes t) e_{t,t}$$

Because $b \rtimes g \mapsto A_{b\rtimes g}$ represents the identity, the same is true for $b \rtimes g \mapsto TA_{b\rtimes g}T^{-1}$, see [2, Proposition 5.1.2]. □

Example 5.2.4. We give an example to show that the adjointness between of τ and $\rtimes G$ of Theorem 5.2.1 fails to hold at the algebra level. Let $G = \mathbb{Z}_2 = \{1, \sigma\}$, $A = \ell$ and $B = (\ell G)^*$ the dual algebra of ℓG with the regular action. Note $\hom_{G\text{-Alg}}(A^\tau, B)$ has two elements only:

$$\varphi_i : \ell \to (\ell G)^* \qquad \varphi_0(1) = 0 \qquad \varphi_1(1) = \chi_1 + \chi_\sigma$$

One the other hand $\hom_{\mathrm{Alg}}(A, B \rtimes G) = \hom_{\mathrm{Alg}}(\ell, (\ell G)^* \rtimes G)$ has at least as many elements as ℓ. For each $\lambda \in \ell$ we can define

$$\varphi_\lambda : \ell \to (\ell G)^* \rtimes G \quad \varphi_\lambda(1) = \chi_1 \rtimes 1 + \lambda(\chi_1 \rtimes \sigma) \quad \lambda \in \ell$$

Note φ_λ is an algebra morphism because $\chi_1 \rtimes 1 + \lambda(\chi_1 \rtimes \sigma)$ is an idempotent element.

Write ψ_{GJ} for the isomorphism of the Theorem 5.2.1

$$\psi_{GJ} : \mathrm{kk}^G(B^\tau, A) \to \mathrm{kk}(B, A \rtimes G) \qquad \psi_{GJ} = \alpha^* \circ \rtimes G \tag{5.2.5}$$

where α is the morphism defined in (5.2.2).

Corollary 5.2.6. *Let G be a finite group such that $1/|G| \in \ell$. Let A be a G-algebra, then*

$$\mathrm{kk}^G(\ell, A) \simeq \mathrm{kk}(\ell, A \rtimes G) \simeq \mathrm{KH}(A \rtimes G)$$

□

6. Induction and Restriction

In this section we study the adjoitness property of the functors of induction and restriction between $\mathfrak{KK}^H$ and $\mathfrak{KK}^G$ where G is a group and H is a subgroup of G.

Let A be a G-algebra and $H \subset G$ a subgroup. If we restrict the action to H we obtain an H-algebra $\mathrm{Res}^{\mathrm{H}}_{\mathrm{G}}(A)$. It is clear this construction defines a functor

$\mathrm{Res}^{\mathrm{H}}_{\mathrm{G}} : G\text{-Alg} \to H\text{-Alg}$. It is easily seen that we can extend $\mathrm{Res}^{\mathrm{H}}_{\mathrm{G}} : G\text{-Alg} \to H\text{-Alg}$ to a triangle functor $\mathrm{Res}^{\mathrm{H}}_{\mathrm{G}} : \mathfrak{KK}^G \to \mathfrak{KK}^H$ so that the following diagram commutes

$$\begin{array}{ccc} G\text{-Alg} & \xrightarrow{\mathrm{Res}^{\mathrm{H}}_{\mathrm{G}}} & H\text{-Alg} \\ {\scriptstyle j^G}\downarrow & & \downarrow{\scriptstyle j^H} \\ \mathfrak{KK}^G & \xrightarrow[\mathrm{Res}^{\mathrm{H}}_{\mathrm{G}}]{} & \mathfrak{KK}^H \end{array}.$$

Let $\pi : G \to G/H$ the projection and A an H-algebra. Consider

$$A^{(G,H)} := \{f : G \to A : \#\pi(\mathrm{supp}(f)) < \infty\}$$

and define

$$\mathrm{Ind}^{\mathrm{G}}_{\mathrm{H}}(A) = \{f \in A^{(G,H)} : \quad f(s) = h(f(sh)) \quad \forall s \in G, \quad h \in H\}.$$

One checks that $\mathrm{Ind}^{\mathrm{G}}_{\mathrm{H}}(A)$ is a G-algebra with pointwise multiplication and the following action of G

$$(g \cdot f)(s) = f(g^{-1}s) \quad f \in \mathrm{Ind}^{\mathrm{G}}_{\mathrm{H}}(A) \quad g, s \in G. \tag{6.1}$$

Observe this construction is functorial, if $\varphi : A \to B$ is a morphism of H-algebras then $\mathrm{Ind}^{\mathrm{G}}_{\mathrm{H}}(\varphi)(f) = \varphi \circ f$.

If $g \in G$, write $\chi_g : G \to \mathbb{Z}$ for the characteristic function. If $a \in A$ and $g \in G$, define

$$\xi_H(g, a) = \sum_{h \in H} \chi_{gh} h^{-1}(a) \qquad \xi_H(g, a)(s) = \begin{cases} h^{-1}(a) & s = gh \\ 0 & s \notin gH \end{cases} \tag{6.2}$$

It is easy to check that these elements belong to $\mathrm{Ind}^{\mathrm{G}}_{\mathrm{H}}(A)$ and every element $\phi \in \mathrm{Ind}^{\mathrm{G}}_{\mathrm{H}}(A)$ can be written as a finite sum

$$\phi = \sum_{g \in \mathcal{R}} \xi_H(g, \phi(g)) \tag{6.3}$$

where $r : G/H \to G$ is a pointed section and $\mathcal{R} = r(G/H)$. Note that we have the following relations

$$s \cdot \xi_H(g, a) = \xi_H(sg, a) \tag{6.4}$$

$$\xi_H(g, a)\xi_H(\tilde{g}, \tilde{a}) = \begin{cases} \xi_H(\tilde{g}, \tilde{g}^{-1}g(a)\tilde{a}) & \tilde{g}^{-1}g \in H \\ 0 & \tilde{g}^{-1}g \notin H \end{cases} \tag{6.5}$$

$$\xi_H(g, a) = \xi_H(gh, h^{-1} \cdot a) \quad h \in H \tag{6.6}$$

Proposition 6.7. *Let A be a G-algebra and B be an H-algebra, then*

$$\mathrm{Ind}^{\mathrm{G}}_{\mathrm{H}}(B \otimes \mathrm{Res}^{\mathrm{H}}_{\mathrm{G}} A) \simeq \mathrm{Ind}^{\mathrm{G}}_{\mathrm{H}}(B) \otimes A$$

Proof. The isomorphisms are given by

$$\begin{array}{rccc} S: & \mathrm{Ind}^{\mathrm{G}}_{\mathrm{H}}(B) \otimes A & \to & \mathrm{Ind}^{\mathrm{G}}_{\mathrm{H}}(B \otimes \mathrm{Res}^{\mathrm{H}}_{\mathrm{G}}(A)) \\ & \xi_H(g, b) \otimes a & \mapsto & \xi_H(g, b \otimes g^{-1} \cdot a) \end{array}$$

$$\begin{array}{rccc} T: & \mathrm{Ind}^{\mathrm{G}}_{\mathrm{H}}(B \otimes \mathrm{Res}^{\mathrm{H}}_{\mathrm{G}} A) & \to & \mathrm{Ind}^{\mathrm{G}}_{\mathrm{H}}(B) \otimes A \\ & \xi_H(g, b \otimes a) & \mapsto & \xi_H(g, b) \otimes g \cdot a \end{array}$$

□

Corollary 6.8. *Let A be a G-algebra. Then*

$$\mathrm{Ind}_{\mathrm{H}}^{\mathrm{G}}\mathrm{Res}_{\mathrm{G}}^{\mathrm{H}}A \to \ell^{(G/H)}\otimes A \qquad \xi_H(s,b)\mapsto \chi_{sH}\otimes s\cdot b$$

is an isomorphism of G-algebras.

□

Proposition 6.9. *Let* $\mathrm{Ind}: H\text{-Alg}\to G\text{-Alg}$ *be the following functor*

$$\mathrm{Ind}(A)=\mathrm{Ind}_{\mathrm{H}}^{\mathrm{G}}(M_H\otimes A).$$

There exists a functor $\mathrm{Ind}_{\mathrm{H}}^{\mathrm{G}}:\mathfrak{KK}^H\to\mathfrak{KK}^G$ *such that the following diagram is commutative*

$$\begin{array}{ccc} H\text{-Alg} & \xrightarrow{\ \mathrm{Ind}\ } & G\text{-Alg} \\ {\scriptstyle j^H}\downarrow & & \downarrow{\scriptstyle j^G} \\ \mathfrak{KK}^H & \xrightarrow[\ \mathrm{Ind}_{\mathrm{H}}^{\mathrm{G}}\]{} & \mathfrak{KK}^G \end{array} \tag{6.10}$$

Proof. Straightforward. □

Proposition 6.11. *The functor* $\mathrm{Ind}_{\mathrm{H}}^{\mathrm{G}}:\mathfrak{KK}^H\to\mathfrak{KK}^G$ *is a triangle functor.*

Proof. Let $f:A\to B$ be a morphism of H-algebras. The following is an isomorphism of G-algebras

$$\Theta:\mathrm{Ind}_{\mathrm{H}}^{\mathrm{G}}(P_f)\to P_{\mathrm{Ind}_{\mathrm{H}}^{\mathrm{G}}(f)} \qquad \xi_H(g,(tp(t),a))\mapsto(\xi_H(g,tp(t)),\xi_H(g,a)). \tag{6.12}$$

The image of the induction functor applied to the path extension of f is

$$\mathrm{Ind}_{\mathrm{H}}^{\mathrm{G}}(\Omega B)\xrightarrow{\mathrm{Ind}_{\mathrm{H}}^{\mathrm{G}}\iota}\mathrm{Ind}_{\mathrm{H}}^{\mathrm{G}}(P_f)\xrightarrow{\mathrm{Ind}_{\mathrm{H}}^{\mathrm{G}}\pi_f}\mathrm{Ind}_{\mathrm{H}}^{\mathrm{G}}A\xrightarrow{\mathrm{Ind}_{\mathrm{H}}^{\mathrm{G}}f}\mathrm{Ind}_{\mathrm{H}}^{\mathrm{G}}B \tag{6.13}$$

By Proposition 6.7 and (6.12), the extension (6.13) is isomorphic to

$$\Omega\,\mathrm{Ind}_{\mathrm{H}}^{\mathrm{G}}(B)\xrightarrow{\mathrm{Ind}_{\mathrm{H}}^{\mathrm{G}}\iota}P_{\mathrm{Ind}_{\mathrm{H}}^{\mathrm{G}}f}\xrightarrow{\pi_{\mathrm{Ind}_{\mathrm{H}}^{\mathrm{G}}f}}\mathrm{Ind}_{\mathrm{H}}^{\mathrm{G}}A\xrightarrow{\mathrm{Ind}_{\mathrm{H}}^{\mathrm{G}}f}\mathrm{Ind}_{\mathrm{H}}^{\mathrm{G}}B.$$

Then $\mathrm{Ind}_{\mathrm{H}}^{\mathrm{G}}:\mathfrak{KK}^H\to\mathfrak{KK}^G$ is a triangle functor. □

Theorem 6.14. *Let G be a group and H a subgroup of G. Then the functors*

$$\mathrm{Ind}_{\mathrm{H}}^{\mathrm{G}}:\mathfrak{KK}^H\to\mathfrak{KK}^G \qquad \mathrm{Res}_{\mathrm{G}}^{\mathrm{H}}:\mathfrak{KK}^G\to\mathfrak{KK}^H$$

are adjoint. Hence

$$\mathrm{kk}^G(\mathrm{Ind}_{\mathrm{H}}^{\mathrm{G}}(B),A)\simeq\mathrm{kk}^H(B,\mathrm{Res}_{\mathrm{G}}^{\mathrm{H}}(A)) \qquad \forall B\in H\text{-Alg}\quad A\in G\text{-Alg}$$

Proof. Let $A\in G$-Alg and $B\in H$-Alg. We need natural transformations

$$\alpha_A\in\mathrm{kk}^G(\mathrm{Ind}_{\mathrm{H}}^{\mathrm{G}}\mathrm{Res}_{\mathrm{G}}^{\mathrm{H}}A,A) \qquad \beta_B\in\mathrm{kk}^H(B,\mathrm{Res}_{\mathrm{G}}^{\mathrm{H}}\mathrm{Ind}_{\mathrm{H}}^{\mathrm{G}}B)$$

which verify the unit and counit condition, respectively.

Define $\varphi_A:\mathrm{Ind}_{\mathrm{H}}^{\mathrm{G}}(\mathrm{Res}_{\mathrm{G}}^{\mathrm{H}}(A))\to M_{G/H}\otimes A$ such that

$$\varphi_A(\xi_H(s,b))=e_{sH,sH}\otimes s\cdot b.$$

One checks that φ_A is a G-equivariant algebra morphism. Put

$$\psi_B:B\to\mathrm{Res}_{\mathrm{G}}^{\mathrm{H}}\mathrm{Ind}_{\mathrm{H}}^{\mathrm{G}}(B)\quad\psi_B(b)=\xi_H(e,b) \tag{6.15}$$

It is easy to check that ψ_B is well-defined and is a map of H-algebras. Let $\alpha_A\in\mathrm{kk}^G(\mathrm{Ind}_{\mathrm{H}}^{\mathrm{G}}\mathrm{Res}_{\mathrm{G}}^{\mathrm{H}}A,A)$ the element represented by φ_A and $\beta_B\in\mathrm{kk}^H(B,\mathrm{Res}_{\mathrm{G}}^{\mathrm{H}}\mathrm{Ind}_{\mathrm{H}}^{\mathrm{G}}B)$ the element represented by ψ_B.

The composite $\mathrm{Res}^{\mathrm{H}}_{\mathrm{G}}(\alpha_A)\circ\beta_{\mathrm{Res}^{\mathrm{H}}_{\mathrm{G}}A}$ is represented by $\mathrm{Res}^{\mathrm{H}}_{\mathrm{G}}(\varphi_A)\circ\psi_{\mathrm{Res}^{\mathrm{H}}_{\mathrm{G}}A}$ which is kk^H-equivalent to the identity in the sense of remark 2.6.2. The element $\alpha_{\mathrm{Ind}^{\mathrm{G}}_{\mathrm{H}}B}\circ \mathrm{Ind}^{\mathrm{G}}_{\mathrm{H}}(\beta_B)\in \mathrm{kk}^G(\mathrm{Ind}^{\mathrm{G}}_{\mathrm{H}}B, \mathrm{Ind}^{\mathrm{G}}_{\mathrm{H}}B)$ is represented by

$$\gamma : \mathrm{Ind}^{\mathrm{G}}_{\mathrm{H}}B \to M_{G/H}\otimes \mathrm{Ind}^{\mathrm{G}}_{\mathrm{H}}B \qquad \xi_H(g,b)\mapsto e_{gH,gH}\otimes \xi_H(g,b)$$

The following morphism of H-algebras

$$\theta : C\to M_{G/H}\otimes C \qquad \theta(c)=e_{H,H}\otimes c$$

represents to the identity in the sense of remark 2.6.2. Then $\mathrm{Ind}^{\mathrm{G}}_{\mathrm{H}}(\theta)$ is kk^G-equivalent to the identity. It is easy to check $\mathrm{Ind}^{\mathrm{G}}_{\mathrm{H}}(\theta)=\gamma$ with $C=B$. □

Write ψ_{IR} for the isomorphism

$$\psi_{IR} : \mathrm{kk}^G(\mathrm{Ind}^{\mathrm{G}}_{\mathrm{H}}B, A)\to \mathrm{kk}^H(B, \mathrm{Res}^{\mathrm{H}}_{\mathrm{G}}A) \qquad \psi_{IR}=\psi_B{}^*\circ \mathrm{Res}^{\mathrm{H}}_{\mathrm{G}} \tag{6.16}$$

where ψ_B is the morphism defined (6.15).

Corollary 6.17. *Let G be a group, H a finite subgroup of G and A a G-algebra then*

$$\mathrm{kk}^G(\ell^{(G/H)}, A)\simeq \mathrm{kk}(\ell, A\rtimes H)\simeq \mathrm{KH}(A\rtimes H)$$

Proof. The isomorphism is the composition of ψ_{GJ} and ψ_{IR} defined in (5.2.5) and in (6.16). □

7. Baaj-Skandalis Duality

In this section we define crossed product functors between the categories G-Alg and G_{gr}-Alg. We prove that they extend to equivalences between $\mathfrak{KK}^G$ and $\hat{\mathfrak{KK}}^G$. In this way we obtain an algebraic duality theorem similar to the duality given by Baaj-Skandalis in [1].

Let A be a G-algebra. Then

$$A\rtimes G=\bigoplus_{s\in G} A\rtimes s \qquad \text{and } (A\rtimes s)(A\rtimes t)\subset A\rtimes st$$

thus $A\rtimes G$ is a G-graded algebra. If $f : A\to B$ is a homomorphism of G-algebras then $f\rtimes G : A\rtimes G\to B\rtimes G$ is a graded homomorphism. Hence we have a functor

$$\rtimes G : G\text{-Alg}\to G_{gr}\text{-Alg}$$

We can also define a functor

$$G\hat{\rtimes} : G_{gr}\text{-Alg}\to G\text{-Alg}$$

as follows. Let B be a G-graded algebra. Let $G\hat{\rtimes}B$ be the algebra which as a module is $\ell^{(G)}\otimes B$ and the product is the following

$$(\chi_g\rtimes a)(\chi_h\rtimes b):=\chi_g\rtimes a_{g^{-1}h}b. \tag{7.1}$$

Here b_g is the homogeneous element associated to g in the decomposition

$$b=\sum_{g\in G} b_g.$$

One checks that the product (7.1) is associative and the action of G, $s \cdot \chi_g \rtimes a = \chi_{sg} \rtimes a$, makes it into a G-algebra. The crossed product $G\hat{\rtimes}B \rtimes G$ is a G-graded algebra which contains B as a graded subalgebra by

$$b \mapsto \sum_{g\in G} \chi_{1_G} \rtimes b_g \rtimes g. \tag{7.2}$$

In (3.2.1) we defined $M_G B$ a G-graded matrix algebra asociated to B. Through the inclusion $b \mapsto e_{1_G,1_G} \otimes b$ we can see B as a G-graded subalgebra of it. In Proposition 7.4 below, we prove that $G\hat{\rtimes}B \rtimes G$ is isomorphic to $M_G B$ as a G-graded algebras.

If $f : A \to B$ is a homogeneous homomorphism we define a G-algebra homomorphism $(G\hat{\rtimes}f)$ in the obvious way. Thus we have a functor

$$\hat{\rtimes}G : G_{gr}\text{-Alg} \to G\text{-Alg}. \tag{7.3}$$

Proposition 7.4. *Let A be a G-algebra and let B be a G-graded algebra.*

a) *There are natural isomorphisms of G-algebras*

$$G\hat{\rtimes}(A \rtimes G) \simeq M_G \otimes A$$

b) *There are natural isomorphisms of G-graded algebras*

$$(G\hat{\rtimes}B) \rtimes G \simeq M_G \otimes B$$

Proof. a) Define $T : G\hat{\rtimes}(A \rtimes G) \to M_G \otimes A$ as

$$T(\chi_g \rtimes a \rtimes s) = g \cdot a \otimes e_{g,gs}.$$

It is easy to check that T is an equivariant algebra isomorphism with inverse given by

$$S(a \otimes e_{r,t}) := \chi_r \rtimes r^{-1} \cdot a \rtimes r^{-1}t$$

b) Define $T : (G\hat{\rtimes}B) \rtimes G \to M_G \otimes B$ as

$$T(\chi_h \rtimes b \rtimes s) = \sum_{r\in G} e_{h,s^{-1}hr} \otimes b_r.$$

It is easy to check that T is a graded algebra isomorphism with inverse given by

$$S(e_{r,s} \otimes b_q) = \chi_r \rtimes b_q \rtimes rqs^{-1} \tag{7.5}$$

□

Theorem 7.6. *The functors $\rtimes G$ and $G\hat{\rtimes}$ extend to inverse equivalences*

$$- \rtimes G : \mathfrak{KK}^G \longrightarrow \hat{\mathfrak{KK}}^G \qquad G\hat{\rtimes} - : \hat{\mathfrak{KK}}^G \longrightarrow \mathfrak{KK}^G$$

Hence if A and B are G-algebras and C and D are G-graded algebras then

$$\mathrm{kk}^G(A,B) \simeq \hat{\mathrm{kk}}^G(A \rtimes G, B \rtimes G) \quad \hat{\mathrm{kk}}^G(C,D) \simeq \mathrm{kk}^G(G\hat{\rtimes}C, G\hat{\rtimes}D)$$

Proof. As $\rtimes G$ maps split sequences to split sequences, $j^G(- \rtimes G)$ is excisive. By Proposition 5.1.2 $j^G(- \rtimes G)$ is G-stable and homotopy invariant, whence it extends to $- \rtimes G : \mathfrak{KK}^G \to \hat{\mathfrak{KK}}^G$ by universality. Similary, as $G\hat{\rtimes}$ maps split exact sequences to split exact sequences then $j^G(G\hat{\rtimes}-)$ is excisive. Because $G\hat{\rtimes}$ maps graded homotopies to equivariant homotopies and $j^G(G\hat{\rtimes}-)$ is M_∞-stable, $j^G(G\hat{\rtimes}-)$ extends to $G\hat{\rtimes} - : \hat{\mathfrak{KK}} \to \mathfrak{KK}^G$ by universality. To finish we must show that the maps

$$\mathrm{kk}^G(A,B) \to \hat{\mathrm{kk}}^G(A \rtimes G, B \rtimes G) \quad \text{and} \quad \hat{\mathrm{kk}}^G(C,D) \to \mathrm{kk}^G(G\hat{\rtimes}C, G\hat{\rtimes}D)$$

are isomorphisms. This is true by Proposition 7.4. □

References

[1] S. Baaj and G. Skandalis, C^*-algèbres de Hopf et théorie de Kasparov équivariante (Hopf C^*-algebras and equivariant Kasparov theory). *K-Theory, 2 (1989), no. 6, 683-721.*

[2] G. Cortiñas and A. Thom. Bivariant algebraic K-theory. *J. Reine und Angew. Math. 610 (2007) 71-123.*

[3] J. Cuntz. A new look at KK-theory. *K-theory 1 (1987) no. 1, 31-51.*

[4] J. Cuntz. Bivariant K-theory and the Weyl algebra. *K-theory 35 (2005), no. 1-2, 93-137.*

[5] P.G. Goerss and J.F. Jardine. Simplicial homotopy theory. *Progress in Mathematics, 174. Birkhäuser Verlag, Basel, 1999.*

[6] E. Guentner, N. Higson and J Trout. Equivariant E-theory for C^*-algebras. *Mem. Amer. Math. Soc. 148 (2000), no. 703.*

[7] N. Higson. A characterization of KK-theory. *Pacific J. Math. 126 (1987), no. 2, 253-276.*

[8] G.G. Kasparov The operator K-functor and extensions of C^*-algebras. *Izv. Akad. Nauk. SSR Ser. Mat. 44 (1980), no. 3, 571-636, 719.*

[9] B. Keller Derived categories and their uses. *Handbook of algebra, Vol. 1, 671701, North-Holland, Amsterdam, 1996*

[10] S. Mac Lane. Categories for the working mathematician. *Second edition. Graduate Texts in Mathematics, 5. Springer-Verlag, New York, 1998.*

[11] R. Meyer. Categorical aspects of bivariant K-theory. *K-theory and noncommutative geometry, 1-39, EMS Ser. Congr. Rep., Eur. Math. Soc., Zürich, 2008.*

[12] C. Weibel. Homotopy Algebraic K-theory. Algebraic K-theory and algebraic number theory (Honolulu, HI, 1987), 461-488, Contemp. Math., 83, Amer. Math. Soc., Providence, RI, 1989.

E-mail address: eugenia@cmat.edu.uy - eugenia.ellis@fu-berlin.de

CMAT, Facultad de Ciencias-UDELAR, Iguá 4225, 11400 Montevideo, Uruguay
Institut für Mathematik, FU, Arnimallee 7, Raum 212, 14195 Berlin, Germany.